

-continued

Cys Ser Asp Leu Glu Val Tyr Phe Cys His Leu Asp Ile Ile Trp  
 1 5 10 15

## (2) INFORMATION FOR SEQ ID NO:17:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (ix) FEATURE:

- (A) NAME/KEY: Disulfide-bond  
 (B) LOCATION: 1..9

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Cys Ser Asn Lys Gln Val Tyr Phe Cys His Leu Asp Ile Ile Trp  
 1 5 10 15

## (2) INFORMATION FOR SEQ ID NO:18:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (ix) FEATURE:


- (A) NAME/KEY: Disulfide-bond  
 (B) LOCATION: 4..8

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:


Asp Lys Glu Cys Val Tyr Phe Cys His Leu Asp Ile Ile Trp  
 1 5 10

We claim:

## 1. A compound of formula I




40



I

wherein

AA<sub>1</sub> is

wherein

X is

hydrogen,  
 alkyl,  
 alkenyl,  
 alkynyl,  
 cycloalkyl,  
 heterocycloalkyl,  
 aryl, or  
 heteroaryl,


n is zero or an integer of 1 or 2,

R<sup>3</sup> is -S-R<sup>4</sup>, wherein R<sup>4</sup> is as defined hereinafter,

45

wherein R<sup>4</sup> is as defined hereinafter,  
 -C(CH<sub>3</sub>)<sub>2</sub>-S-R<sup>4</sup>, wherein R<sup>4</sup> is covalently linked  
 to AA<sup>1a</sup> through a disulfide bridge,


50



55

wherein X is as defined above, X<sup>1</sup> is hydrogen or  
 Asp-Lys-Glu and n and R<sup>3</sup> are as defined above,

60



65

wherein X, n, and R<sup>3</sup> are as defined above, or